

Bioprocessing of agri-food residues into value-added products

ShT-02.1-3

E. Maresca^I, M. Aulitto^I, A. Alfano^{II}, R. Ausiello^I, D. Cimini^{III}, C. Schiraldi^{II}, P. Contursi^I

^IUniversità Federico II di Napoli, Dpt Biologia, Complesso Universitario di Montesantangelo, NAPOLI, Italy, ^{II}Università della Campania Luigi Vanvitelli, Dpt. di Medicina Sperimentale, Via de Crecchio 7, NAPOLI, Italy, ^{III}Dpt di Scienze e Tecnologie Ambientali Biologiche e Farmaceutiche, Università degli Studi della Campania Luigi Vanvitelli, Via Vivaldi 43, Caserta, Italy

Sustainable and resilient productions are based on a circular economy and minimal processing. In this frame, the valorization of agri-food residues (AFR) is a hot topic of research in zero-waste global development and for the green production of biomaterials such as those based on bacterial cellulose (BC) and lactic acid (LA). Among AFR, citrus residues have been successfully used as carbon sources in culture media for microbial fermentation (1). However, existing methods for valorizing citrus waste often require chemical pretreatments of the biomass, which have a high environmental impact (2). In order to establish an ecofriendly and sustainable strategy for production of LA and BC we pointed to a moderately thermophilic lactic acid producer *Weizmannia coagulans* (3) strain and to a bacterial/yeast community (Kombucha) respectively, exploiting their degradative ability of complex biomasses. Our result show: i) the feasibility of replacing the standard growth medium with less expensive carbon/nitrogen which are widely available at low-cost, and suitable for fermentation growth; ii) the setting up of a green process for BC and LA production based on untreated biomass; iii) the potential of citrus waste to release bioactive compounds with several health-promoting and therapeutic advantages upon microbial fermentation.

1. Aulitto M, Alfano A, Maresca E, et al. Thermophilic biocatalysts for one-step conversion of citrus waste into lactic acid. *Appl Microbiol Biotechnol.* 2024;108(1):155
2. Huang, R et al. Enhanced ethanol production from pomelo peel waste by integrated hydrothermal treatment, multienzyme formulation, and fed-batch operation. *Journal of agricultural and food chemistry.* 2014; 62(20): 4643-51
3. Aulitto M, Fusco S, Bartolucci S, et al. “*Bacillus coagulans* MA-13: a promising thermophilic and cellulolytic strain for the production of lactic acid from lignocellulosic hydrolysate.” *Biotechnology for biofuels.* 2017; 10: 210